

Page 1 of 19 05/30/2011

How often the home health team taught patients (or their family caregivers) about their drugs

85%

Page 2 of 19 05/30/2011

How often the home health team checked patients' risk of falling

94%

Page 3 of 19 05/30/2011

How often the home health team checked patients for depression

93%

Page 4 of 19 05/30/2011

How often the home health team determined whether patients received a flu shot for the current flu season

63%

Page 5 of 19 05/30/2011

How often the home health team determined whether their patients received a pneumococcal vaccine (pneumonia shot)

60%

Page 6 of 19 05/30/2011

With diabetes, how often the home health team got doctor's orders, gave foot care, and taught patients about foot care

85%

Page 7 of 19 05/30/2011

How often the home health team checked patients for pain



Page 8 of 19 05/30/2011

How often the home health team treated their patients' pain

95%

Page 9 of 19 05/30/2011

How often the home health team treated heart failure (weakening of the heart) patients' symptoms



Page 10 of 19 05/30/2011

How often the home health team took doctor-ordered action to prevent pressure sores (bed sores)

89%

Page 11 of 19 05/30/2011

How often the home health team included treatments to prevent pressure sores (bed sores) in the plan of care


90%

Page 12 of 19 05/30/2011


How often the home health team checked patients for the risk of developing pressure sores (bed sores)

96%


Page 13 of 19 05/30/2011



Page 14 of 19 05/30/2011



Page 15 of 19 05/30/2011



Page 16 of 19 05/30/2011



Page 17 of 19 05/30/2011

How often patients receiving home health care needed urgent, unplanned care in the ER without being admitted

Page 18 of 19 05/30/2011

How often home health patients had to be admitted to the hospital

29%

Page 19 of 19 05/30/2011